

Using height association studies to gain insights into human idiopathic short and syndromic stature phenotypes

Guillaume Lettre

Received: 16 April 2012 / Revised: 5 June 2012 / Accepted: 29 June 2012 / Published online: 2 September 2012
© IPNA 2012

Abstract Variation in adult height is not the most clinically relevant human quantitative trait, yet its study provides the foundation of many quantitative genetics theories and important statistical concepts (e.g. regression). Even today, the analysis of adult height by genome-wide association studies (GWAS) continues to significantly impact human genetics: these studies have led to the discovery of >200 loci associated with variation in adult height and have highlighted the very polygenic nature of human continuous traits. In this brief review, I discuss and provide examples on how such genetic associations, identified in individuals of *normal* height, could help understand the complex genetics behind such phenotypes as idiopathic short stature (ISS) or extreme/syndromic height phenotypes of unknown cause.

Keywords Adult height · Stature · Idiopathic short stature · Syndromic height · Genome-wide association study · Genetics

Introduction

Childhood growth is one of the main traits monitored by pediatricians around the world to ensure that a child's development is normal. Although the study of height variation in humans has been at the center stage of quantitative genetics since the birth of this discipline [1–3], surprisingly little is known about the factors that regulate developmental growth, especially in terms of explaining the interindividual variation in height observed among children of the same age. The environment undoubtedly plays an important role

(e.g. nutrition, infections), but it is also clear that genetic programs influence birth length, puberty growth spurt, growth velocity and final adult height [4–6]. However, it is only with the emergence of new genomic technologies that the first genetic polymorphisms that influence height were recently identified [7, 8]. These initial discoveries paved the way for larger follow-up studies that have so far yielded >200 single nucleotide polymorphisms (SNPs) convincingly associated with height variation in adults [9]. In this review, I focus on how such genetic discoveries, which help understand variation in height observed in the general population, could also help to elucidate the genetic basis of *abnormal* phenotypes, such as idiopathic short stature (ISS) or extreme short/tall height, caused by mutations in unknown genes.

Genetic association studies of adult height

The recent insights into adult height variation through genome-wide association studies (GWAS) have extensively been reviewed [10–12]. Therefore, I will briefly summarize here only the main findings that will help guide our discussion. In 2010, the Genetic Investigation of ANthropometric Traits (GIANT) Consortium (an international consortium that gathers >50 studies with GWAS results; http://www.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium) published what is still to date one of the largest meta-analysis of genome-wide association results for a human continuous trait: the GIANT investigators combined results from >180,000 participants and identified 180 SNPs associated with adult height at genome-wide significance [9]. Importantly, this study confirmed the polygenic architecture of height—which is that the segregation of hundreds of genetic variants, each with a small effect on height, is sufficient to account for the variation in stature among human populations. The 180 GIANT SNPs together explain approximately 10 % of the phenotypic

G. Lettre (✉)
Montreal Heart Institute and Université de Montréal,
5000 Bélanger Street,
Montreal, Quebec H1T 1C8, Canada
e-mail: guillaume.lettre@mhi-humangenetics.org

variation in adult height. Besides the logistical *tour de force* of the study itself, the GIANT study identified several genomic regions and biological pathways previously unsuspected to play a role in human height (Table 1). For most of these loci, two important questions remain: (1) which genes play a role in human stature and (2) how genetic variation within these loci influences the functions of these genes? The few examples of successful functional characterization of GWAS hits suggest that the translation from genetic results to biological insights requires an integrative approach that considers genetic and genomic datasets, as well as molecular and cell biology experiments and characterization in animal models [13–16].

The GIANT study analyzed exclusively Europeans or individuals of European ancestry [9], but large meta-analyses of GWAS have also analyzed adult height in other ethnic groups. For example, the KARE Consortium analyzed height in 8,842 South Koreans and found 15 loci associated with height, many of which had already been reported by the GIANT study [17, 18]. Two large association studies were recently performed in African-ancestry individuals, and both reported a limited number of associations not previously identified in populations of European ancestry. Furthermore, both studies also replicated most associations that were initially reported by the GIANT Consortium [19, 20]. Although population-specific associations were observed for this trait, these results suggest that most height-associated genetic polymorphisms have *trans*-ethnic effect.

Genetic associations during childhood growth

Much less is known about the genetic factors that modulate growth during childhood despite high heritability estimates [21]. This lack of information is attributed partly to the heterogeneity of growth timing and velocity, even in children of

the same sex and age. Therefore, the study of the genetics of human growth requires longitudinal height measures obtained over several years (ideally from birth to the end of growth at 18–21 years of age). Fortunately, such large longitudinal cohorts do exist, and attempts have been made to identify genetic determinants of growth, focusing on birth length, growth rate at different periods during development and timing of puberty [22–25]. These studies have confirmed that the SNPs associated with adult height do play an important role in childhood growth. A genetic score based on genotypes at the 180 height SNPs identified by the GIANT Consortium associates strongly with birth length and growth rate [25]. The results of a study in twins suggest that the SNPs associated with adult height affect more prepubertal height than the pubertal growth spurt [24]. Common DNA polymorphisms (SNPs) at the *LIN28B* locus, however, associate with growth velocity, and the effect was proposed by the respective authors to be sex-specific [22, 24]. It is likely that combining genome-wide association results for growth through meta-analyses, as has been done for stature, would yield novel associations because of increased statistical power.

ISS, one of the main reasons for referral to pediatric endocrinology clinics, is defined as a sex-, age- and population-adjusted height that is more than two standard deviations (SD) below the mean with no evidence of endocrine, nutritional or chromosomal anomalies [26]. Understanding the causes of ISS is critical in the context of our fragile healthcare system, given the costs associated with growth hormone therapy (>\$15,000/child/year) [27]. It would be extremely useful for pediatric endocrinologists to be able to determine whether a child with ISS is in the *normal* range of his/her height potential given his/her genetic code and the segregation of common height-associated alleles, or whether something else (e.g. environmental factors, de novo detrimental mutations) is causing the abnormal height phenotype.

Table 1 Biological pathways (KEGG database) that are enriched ($P < 0.05$) for genes located near (≤ 300 kb) height-associated single nucleotide polymorphisms^a

Biological pathway (KEGG)	Genes
Hedgehog signaling pathway	<i>BMP6, IHH, PTCH1, WNT6, WNT9A, FBXW11, HHIP, WNT10A, WNT3A</i>
Gamma-hexachlorocyclohexane degradation	<i>DHRS1, LOC283871</i>
MAPK signaling pathway	<i>ARRB1, CACNB1, CHUK, FGFR3, FGFR4, GNA12, MKNK2, MEF2C, MAP3K3, MOS, GADD45B, NF1, NFATC4, PPM1A, MAPK9, MAP2K3, RASA2, RPS6KA1, TGFB1, TGFB2, TNF, MAP3K14, RASGRP3</i>
Antigen processing and presentation	<i>HLA-B, HLA-C, HLA-DMA, HLA-DMB, HLADOB, HLA-DQA1, HLA-DQA2, HLA-DQB1, HLA-DRA, HLA-DRB1, HLA-DRB5, LTA, PSME1, PSME2, TAP1, TAP2</i>
TGF- β signaling pathway	<i>AMH, BMP6, ID4, LTBP1, TGFB1, TGFB2, TNF, GDF5, CUL1, NOG</i>
Type II diabetes mellitus	<i>INSR, KCNJ11, PKM2, PRKCD, PRKCZ, MAPK9, ABCC8, TNF, SOCS2</i>
FC epsilon RI signaling pathway	<i>CSF2, IL5, IL13, LYN, PRKCD, MAPK9, MAP2K3, TNF</i>
Folate biosynthesis	<i>ATP13A2</i>
Citrate cycle TCA cycle	<i>CS, PC, PCK2, SDHB, SUCLG2</i>

MAPK, Mitogen-activated protein kinase; TGF- β , transforming growth factor beta; TCA, tricarboxylic acid cycle

^aSee [9] for details

In a first step to address this question, Chan and colleagues analyzed the effect of the 180 common height SNPs on stature in adult individuals with an extreme height phenotype (very short or tall; >2 SD below or above the population mean) [28]. They found that at the extreme tall end of the height distribution, phenotypes were consistent with the segregation of common DNA polymorphisms, each with a weak effect on height. However, at the extreme short end of the distribution, phenotypes were more extreme than what was expected based on the polygenic model; this was even more obvious in individuals with extremely short stature (<2.8 SD below the mean or in other words in the 0.25th percentile for height) [28]. The authors suggest that non-additive genetic or environmental factors are likely to contribute to final height mostly in extremely short individuals (Fig. 1). Validating these findings in large cohorts of ISS children would be extremely valuable in pediatric endocrinology.


Common height variation and Mendelian genetics

Another interesting finding from the GIANT study is that SNPs associated with adult height are found more often near skeletal- or growth defect-related genes than can be expected by chance [9]. This is a strong argument supporting the fact that height loci are not randomly distributed

across the genome but rather clustered near meaningful candidate genes. Furthermore, it indicates that hypomorphic alleles which alter the expression level or the biological activity of genes that are otherwise known to cause severe skeletal dysplasia or extreme height phenotypes when inactivated influence *normal* height variation. Height is not alone in this category of phenotypes, and we now have several combinations of genes–phenotypes for which an allelic spectrum has been linked to variable mild-to-severe phenotypic manifestations (e.g. *PCSK9* and low-density lipoprotein cholesterol/coronary artery disease, *MC4R* and body mass index/obesity). Next-generation DNA re-sequencing of large cohorts, for height and other traits should disentangle loci/gene and determine whether allelic series are a general feature of the genetic architecture of human complex diseases and traits.

The proximity between height-associated SNPs and skeletal growth genes can also be the basis of the following hypothesis: genes located near height SNPs are excellent candidates to underlie more extreme Mendelian skeletal growth phenotypes. A survey of the Online Mendelian Inheritance in Man (OMIM) database with the keywords “overgrowth,” “height,” “stature” or “dysplasia” revealed 316 Mendelian phenotypes or loci with an unknown molecular basis (<http://ncbi.nlm.nih.gov/omim>; accessed 29 Mar 2012). For those phenotypes for which disease loci have been mapped between chromosome bands, one strategy that could possibly be adopted to find the causal genes and

Fig. 1 Possible model of the genetic architecture of height across the phenotypic distribution. Familial and de novo mutations are too rare to account for the variation in height observed in the general population. Common DNA sequence variants [e.g. single nucleotide polymorphisms (SNPs)] are associated with height in the general population as well as in very short or tall individuals [± 2 standard deviations (*SD*)], although some results suggest that they may not cause extreme height phenotypes [28]. The role of rare sequence variants in height variation is currently unknown



mutations would be to sequence the genes within the interval that falls near SNPs associated with height.

This approach has not yet been applied systematically, but there are at least two indirect examples that show its great promises, i.e. the *TRIP11* gene and the *mir-17~92* microRNA locus. In 2008, we and others found SNPs associated with adult height near the *TRIP11* gene [29, 30]. Mutations in *TRIP11* were later found to cause skeletal dysplasia in the mouse and achondroplasia type 1A in humans [31]. More recently, using comparative genomic hybridization arrays, de Pontual and colleagues found that deletions of the microRNA cluster locus *mir-17~92* can cause Feingold syndrome, which is characterized by microcephaly, short stature and digital anomalies [32]. The GIANT Consortium had presented evidence of an association with adult height for a SNP (rs90822575) located only 17.4 kb downstream of *mir-17~92* [9]. The link between microRNAs and human height is intriguing, given the strong association of the *LIN28B* locus with adult height and human growth (see above) and the fact that we found an enrichment of *let-7* target genes among height associated loci [29].

Conclusions

After more than 100 years of research, the study of the genetics of human height continues to be at the forefront of the field of quantitative genetics. Given the progress in DNA sequencing technology and the availability of height measures in very large re-sequenced cohorts, this will continue for years to come. Discoveries from GWAS have highlighted novel biological pathways that influence adult height and also, to some extent, human growth. Although more genetic work is needed, functional studies based on these genetic findings are already changing our understanding of growth in animal models and humans [16]. Finally, SNPs associated with height and their nearby genes could have clinical applications, for example by helping us to understand the etiology of ISS or orphan syndromes with skeletal defects or other growth phenotypes.

Acknowledgments I would like to thank Aikaterini Kritikou and members of my laboratory for comments on earlier versions of this manuscript. Work on the genetics of height and other anthropometric traits in my laboratory is funded by the Montreal Heart Institute Foundation and the “Fonds de Recherche du Québec en Santé (FRQS).”

References

- Galton F (1885) Regression towards mediocrity in hereditary stature. *J R Anthropol Inst* 5:329–348
- Fisher RA (1918) The correlation between relatives on the supposition of Mendelian inheritance. *Trans R Soc Edinb* 52:399–433
- Pearson K, Lee A (1903) On the laws of inheritance in man. I. Inheritance of physical characteristics. *Biometrika* 2:357–462
- Silventoinen K, Haukka J, Dunkel L, Tynelius P, Rasmussen F (2008) Genetics of pubertal timing and its associations with relative weight in childhood and adult height: the Swedish Young Male Twins Study. *Pediatrics* 121:e885–e891
- Perola M, Sarmola S, Hekkinen T, Martin NG, Visscher PM, Montgomery GW, Benyamin B, Harris JR, Boomsma D, Willemsen G, Hottenga JJ, Christensen K, Kyvik KO, Sorensen TI, Pedersen NL, Magnusson PK, Spector TD, Widen E, Silventoinen K, Kaprio J, Palotie A, Peltonen L (2007) Combined genome scans for body stature in 6,602 European twins: evidence for common Caucasian loci. *PLoS Genet* 3:e97
- Visscher PM, Medland SE, Ferreira MA, Morley KI, Zhu G, Cornes BK, Montgomery GW, Martin NG (2006) Assumption-free estimation of heritability from genome-wide identity-by-descent sharing between full siblings. *PLoS Genet* 2:e41
- Weedon MN, Lettre G, Freathy RM, Lindgren CM, Voight BF, Perry JR, Elliott KS, Hackett R, Guiducci C, Shields B, Zeggini E, Lango H, Lyssenko V, Timpson NJ, Burt NP, Rayner NW, Saxena R, Ardlie K, Tobias JH, Ness AR, Ring SM, Palmer CN, Morris AD, Peltonen L, Salomaa V, Smith GD, Groop LC, Hattersley AT, McCarthy MI, Hirschhorn JN, Frayling TM (2007) A common variant of HMG2 is associated with adult and childhood height in the general population. *Nat Genet* 39:1245–1250
- Sanna S, Jackson AU, Nagaraja R, Willer CJ, Chen WM, Bonnycastle LL, Shen H, Timpson N, Lettre G, Usala G, Chines PS, Stringham HM, Scott LJ, Dei M, Lai S, Albai G, Crisponi L, Naitza S, Doheny KF, Pugh EW, Ben-Shlomo Y, Ebrahim S, Lawlor DA, Bergman RN, Watanabe RM, Uda M, Tuomilehto J, Coresh J, Hirschhorn JN, Shuldiner AR, Schlessinger D, Collins FS, Davey Smith G, Boerwinkle E, Cao A, Boehnke M, Abecasis GR, Mohlke KL (2008) Common variants in the GDF5-UQC region are associated with variation in human height. *Nat Genet* 40:198–203
- Lango Allen H, Estrada K, Lettre G, Berndt SI, Weedon MN, Rivadeneira F, Willer CJ, Jackson AU, Vedantam S, Raychaudhuri S, Ferreira T, Wood AR, Weyant RJ, Segre AV, Speliotes EK, Wheeler E, Soranzo N, Park JH, Yang J, Gudbjartsson D, Heard-Costa NL, Randall JC, Qi L, Vernon Smith A, Magi R, Pastinen T, Liang L, Heid IM, Luan J, Thorleifsson G, Winkler TW, Goddard ME, Sin Lo K, Palmer C, Workalemahu T, Aulchenko YS, Johansson A, Zillikens MC, Feitosa MF, Esko T, Johnson T, Ketkar S, Kraft P, Mangino M, Prokopenko I, Absher D, Albrecht E, Ernst F, Glazer NL, Hayward C, Hottenga JJ, Jacobs KB, Knowles JW, Kutalik Z, Monda KL, Polasek O, Preuss M, Rayner NW, Robertson NR, Steinthorsdottir V, Tyrer JP, Voight BF, Wiklund F, Xu J, Zhao JH, Nyholt DR, Pellikka N, Perola M, Perry JR, Surakka I, Tammesoo ML, Altmaier EL, Amin N, Aspelund T, Bhargava T, Boucher G, Chasman DI, Chen C, Coin L, Cooper MN, Dixon AL, Gibson Q, Grundberg E, Hao K, Juhani Junttila M, Kaplan LM, Kettunen J, König IR, Kwan T, Lawrence RW, Levinson DF, Lorentzon M, McKnight B, Morris AP, Muller M, Suh Ngwa J, Purcell S, Rafelt S, Salem RM, Salvi E, Sanna S, Shi J, Sovio U, Thompson JR, Turchin MC, Vandenput L, Verlaan DJ, Vitart V, White CC, Ziegler A, Almgren P, Balmforth AJ, Campbell H, Citterio L, De Grandi A, Dominiczak A, Duan J, Elliott P, Elosua R, Eriksson JG, Freimer NB, Geus EJ, Glorioso N, Haiqing S, Hartikainen AL, Havulinna AS, Hicks AA, Hui J, Igl W, Illig T, Jula A, Kajantie E, Kilpelainen TO, Koivumäki M, Kolcic I, Koskenvuo S, Kovacs P, Laitinen J, Liu J, Lokki ML, Marusic A, Maschio A, Meitinger T, Mulas A, Pare G, Parker AN, Peden JF, Petersmann A, Pichler I, Pietiläinen KH, Pouta A, Ridderstråle M, Rotter JJ, Sambrook JG, Sanders AR, Schmidt CO, Sinisalo J, Smit JH, Stringham HM, Bragi Walters G, Widen E, Wild SH, Willemsen G, Zagato L, Zgaga L, Zitting P, Alavere H, Farrall

- M, McArdle WL, Nelis M, Peters MJ, Ripatti S, van Meurs JB, Aben KK, Ardlie KG, Beckmann JS, Beilby JP, Bergman RN, Bergmann S, Collins FS, Cusi D, den Heijer M, Eiriksdottir G, Gejman PV, Hall AS, Hamsten A, Huikuri HV, Iribarren C, Kahonen M, Kaprio J, Kathiresan S, Kiemeny L, Kocher T, Launer LJ, Lehtimäki T, Melander O, Mosley TH Jr, Musk AW, Nieminen MS, O'Donnell CJ, Ohlsson C, Oostra B, Palmer LJ, Raitakari O, Ridker PM, Rioux JD, Rissanen A, Rivolta C, Schunkert H, Shuldiner AR, Siscovick DS, Stumvoll M, Tonjes A, Tuomilehto J, van Ommen GJ, Viikari J, Heath AC, Martin NG, Montgomery GW, Province MA, Kayser M, Arnold AM, Atwood LD, Boerwinkle E, Chanock SJ, Deloukas P, Gieger C, Gronberg H, Hall P, Hattersley AT, Hengstenberg C, Hoffman W, Lathrop GM, Salomaa V, Schreiber S, Uda M, Waterworth D, Wright AF, Assimes TL, Barroso I, Hofman A, Mohlke KL, Boomsma DI, Caulfield MJ, Cupples LA, Erdmann J, Fox CS, Gudnason V, Gyllenstein U, Harris TB, Hayes RB, Jarvelin MR, Mooser V, Munroe PB, Ouwehand WH, Penninx BW, Pramstaller PP, Quertermous T, Rudan I, Samani NJ, Spector TD, Volzke H, Watkins H, Wilson JF, Groop LC, Haritunians T, Hu FB, Kaplan RC, Metspalu A, North KE, Schlessinger D, Wareham NJ, Hunter DJ, O'Connell JR, Strachan DP, Wichmann HE, Borecki IB, van Duijn CM, Schadt EE, Thorsteinsdottir U, Peltonen L, Uitterlinden AG, Visscher PM, Chatterjee N, Loos RJ, Boehnke M, McCarthy MI, Ingelsson E, Lindgren CM, Abecasis GR, Stefansson K, Frayling TM, Hirschhorn JN (2010) Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature* 467:832–838
10. Perola M (2011) Genetics of human stature: Lessons from genome-wide association studies. *Horm Res Paediatr* 76[Suppl 3]:10–11
 11. Lettre G (2011) Recent progress in the study of the genetics of height. *Hum Genet* 129:465–472
 12. Visscher PM, McEvoy B, Yang J (2010) From Galton to GWAS: quantitative genetics of human height. *Genet Res* 92:371–379
 13. Sankaran VG, Menne TF, Xu J, Akie TE, Lettre G, Van Handel B, Mikkola HK, Hirschhorn JN, Cantor AB, Orkin SH (2008) Human fetal hemoglobin expression is regulated by the developmental stage-specific repressor BCL11A. *Science* 322:1839–1842
 14. Musunuru K, Strong A, Frank-Kamenetsky M, Lee NE, Ahfeldt T, Sachs KV, Li X, Li H, Kuperwasser N, Ruda VM, Pirruccello JP, Muchmore B, Prokunina-Olsson L, Hall JL, Schadt EE, Morales CR, Lund-Katz S, Phillips MC, Wong J, Cantley W, Racie T, Ejebe KG, Orho-Melander M, Melander O, Koteliensky V, Fitzgerald K, Krauss RM, Cowan CA, Kathiresan S, Rader DJ (2010) From noncoding variant to phenotype via SORT1 at the 1p13 cholesterol locus. *Nature* 466:714–719
 15. Genovese G, Friedman DJ, Ross MD, Lecordier L, Uzureau P, Freedman BI, Bowden DW, Langefeld CD, Oleksyk TK, Uscinski Knob AL, Bernhardt AJ, Hicks PJ, Nelson GW, Vanhollenbeke B, Winkler CA, Kopp JB, Pays E, Pollak MR (2010) Association of trypanolytic ApoL1 variants with kidney disease in African Americans. *Science* 329:841–845
 16. Zhu H, Shah S, Shyh-Chang N, Shinoda G, Einhorn WS, Viswanathan SR, Takeuchi A, Grasmann C, Rinn JL, Lopez MF, Hirschhorn JN, Palmert MR, Daley GQ (2010) Lin28a transgenic mice manifest size and puberty phenotypes identified in human genetic association studies. *Nat Genet* 42:626–630
 17. Kim JJ, Lee HI, Park T, Kim K, Lee JE, Cho NH, Shin C, Cho YS, Lee JY, Han BG, Yoo HW, Lee JK (2010) Identification of 15 loci influencing height in a Korean population. *J Hum Genet* 55:27–31
 18. Cho YS, Go MJ, Kim YJ, Heo JY, Oh JH, Ban HJ, Yoon D, Lee MH, Kim DJ, Park M, Cha SH, Kim JW, Han BG, Min H, Ahn Y, Park MS, Han HR, Jang HY, Cho EY, Lee JE, Cho NH, Shin C, Park T, Park JW, Lee JK, Cardon L, Clarke G, McCarthy MI, Lee JY, Oh B, Kim HL (2009) A large-scale genome-wide association study of Asian populations uncovers genetic factors influencing eight quantitative traits. *Nat Genet* 41:527–534
 19. N'Diaye A, Chen GK, Palmer CD, Ge B, Tayo B, Mathias RA, Ding J, Nalls MA, Adeyemo A, Adoue V, Ambrosone CB, Atwood L, Bandera EV, Becker LC, Berndt SI, Bernstein L, Blot WJ, Boerwinkle E, Britton A, Casey G, Chanock SJ, Demerath E, Deming SL, Diver WR, Fox C, Harris TB, Hernandez DG, Hu JJ, Ingles SA, John EM, Johnson C, Keating B, Kittles RA, Kolonel LN, Kritchevsky SB, Le Marchand L, Lohman K, Liu J, Millikan RC, Murphy A, Musani S, Neslund-Dudas C, North KE, Nyante S, Ogunniyi A, Ostrander EA, Papanicolaou G, Patel S, Pettaway CA, Press MF, Redline S, Rodriguez-Gil JL, Rotimi C, Rybicki BA, Salako B, Schreiner PJ, Signorello LB, Singleton AB, Stanford JL, Stram AH, Stram DO, Strom SS, Suktitipat B, Thun MJ, Witte JS, Yanek LR, Ziegler RG, Zheng W, Zhu X, Zmuda JM, Zonderman AB, Evans MK, Liu Y, Becker DM, Cooper RS, Pastinen T, Henderson BE, Hirschhorn JN, Lettre G, Haiman CA (2011) Identification, replication, and fine-mapping of Loci associated with adult height in individuals of African ancestry. *PLoS Genet* 7:e1002298
 20. Carty CL, Johnson NA, Hutter CM, Reiner AP, Peters U, Tang H, Kooperberg C (2012) Genome-wide association study of body height in African Americans: the Women's Health Initiative SNP Health Association Resource (SHARe). *Hum Mol Genet* 21:711–720
 21. Silventoinen K, Pietiläinen KH, Tynelius P, Sorensen TI, Kaprio J, Rasmussen F (2008) Genetic regulation of growth from birth to 18 years of age: the Swedish young male twins study. *Am J Hum Biol* 20:292–298
 22. Widen E, Ripatti S, Cousminer DL, Surakka I, Lappalainen T, Jarvelin MR, Eriksson JG, Raitakari O, Salomaa V, Sovio U, Hartikainen AL, Pouta A, McCarthy MI, Osmond C, Kajantie E, Lehtimäki T, Viikari J, Kahonen M, Tyler-Smith C, Freimer N, Hirschhorn JN, Peltonen L, Palotie A (2010) Distinct variants at LIN28B influence growth in height from birth to adulthood. *Am J Hum Genet* 86:773–782
 23. Sovio U, Bennett AJ, Millwood IY, Molitor J, O'Reilly PF, Timpson NJ, Kaakinen M, Laitinen J, Haukka J, Pillas D, Tzoulaki I, Hoggart C, Coin LJ, Whittaker J, Pouta A, Hartikainen AL, Freimer NB, Widen E, Peltonen L, Elliott P, McCarthy MI, Jarvelin MR (2009) Genetic determinants of height growth assessed longitudinally from infancy to adulthood in the northern Finland birth cohort 1966. *PLoS Genet* 5:e1000409
 24. Vrieze SJ, McGue M, Miller MB, Legrand LN, Schork NJ, Iacono WG (2011) An assessment of the individual and collective effects of variants on height using twins and a developmentally informative study design. *PLoS Genet* 7:e1002413
 25. Paternoster L, Howe LD, Tilling K, Weedon MN, Freathy RM, Frayling TM, Kemp JP, Smith GD, Timpson NJ, Ring SM, Evans DM, Lawlor DA (2011) Adult height variants affect birth length and growth rate in children. *Hum Mol Genet* 20:4069–4075
 26. Ranke MB (2011) Chairman's summary: definition of idiopathic short stature. *Horm Res Paediatr* 76[Suppl 3]:2
 27. Cuttler L, Marinova D, Mercer MB, Connors A, Meehan R, Silvers JB (2009) Patient, physician, and consumer drivers: referrals for short stature and access to specialty drugs. *Med Care* 47:858–865
 28. Chan Y, Holmen OL, Dauber A, Vatten L, Havulinna AS, Skorpén F, Kvaloy K, Silander K, Nguyen TT, Willer C, Boehnke M, Perola M, Palotie A, Salomaa V, Hveem K, Frayling TM, Hirschhorn JN, Weedon MN (2011) Common variants show predicted polygenic effects on height in the tails of the distribution, except in extremely short individuals. *PLoS Genet* 7:e1002439
 29. Lettre G, Jackson AU, Gieger C, Schumacher FR, Berndt SI, Sanna S, Eyheramendy S, Voight BF, Butler JL, Guiducci C, Illig

- T, Hackett R, Heid IM, Jacobs KB, Lyssenko V, Uda M, Boehnke M, Chanock SJ, Groop LC, Hu FB, Isomaa B, Kraft P, Peltonen L, Salomaa V, Schlessinger D, Hunter DJ, Hayes RB, Abecasis GR, Wichmann HE, Mohlke KL, Hirschhorn JN (2008) Identification of ten loci associated with height highlights new biological pathways in human growth. *Nat Genet* 40:584–591
30. Gudbjartsson DF, Walters GB, Thorleifsson G, Stefansson H, Halldorsson BV, Zusmanovich P, Sulem P, Thorlacius S, Gylfason A, Steinberg S, Helgadóttir A, Ingason A, Steinthorsdóttir V, Olafsdóttir EJ, Olafsdóttir GH, Jonsson T, Borch-Johnsen K, Hansen T, Andersen G, Jorgensen T, Pedersen O, Aben KK, Witjes JA, Swinkels DW, den Heijer M, Franke B, Verbeek AL, Becker DM, Yanek LR, Becker LC, Tryggvadóttir L, Rafnar T, Gulcher J, Kiemeneý LA, Kong A, Thorsteinsdóttir U, Stefansson K (2008) Many sequence variants affecting diversity of adult human height. *Nat Genet* 40:609–615
31. Smits P, Bolton AD, Funari V, Hong M, Boyden ED, Lu L, Manning DK, Dwyer ND, Moran JL, Prysak M, Merriman B, Nelson SF, Bonafe L, Superti-Furga A, Ikegawa S, Krakow D, Cohn DH, Kirchhausen T, Warman ML, Beier DR (2010) Lethal skeletal dysplasia in mice and humans lacking the golgin GMAP-210. *N Engl J Med* 362:206–216
32. de Pontual L, Yao E, Callier P, Faivre L, Drouin V, Cariou S, Van Haeringen A, Genevieve D, Goldenberg A, Oufadem M, Manouvrier S, Munnich A, Vidigal JA, Vekemans M, Lyonnet S, Henrion-Caude A, Ventura A, Amiel J (2011) Germline deletion of the miR-17 approximately 92 cluster causes skeletal and growth defects in humans. *Nat Genet* 43:1026–1030